

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C12N 9/42, D21C 9/10</p>	<p>A1</p>	<p>(11) International Publication Number: WO 97/11164 (43) International Publication Date: 27 March 1997 (27.03.97)</p>
<p>(21) International Application Number: PCT/US96/14692 (22) International Filing Date: 13 September 1996 (13.09.96) (30) Priority Data: 530,801 20 September 1995 (20.09.95) US (71) Applicant: GENENCOR INTERNATIONAL, INC. [US/US]; 4 Cambridge Place, 1870 South Winton Road, Rochester, NY 14618 (US). (72) Inventors: BODIE, Elizabeth, A.; Genencor International, Inc., 925 Page Mill Road, Palo Alto, CA 94304-1013 (US). CUEVAS, William, A.; Genencor International, Inc., 925 Page Mill Road, Palo Alto, CA 94304-1013 (US). KANTELINEN, Anne, K.; Genencor International, Inc., 925 Page Mill Road, Palo Alto, CA 94304-1013 (US). (74) Agent: ANDERSON, Kirsten, A.; Genencor International, Inc., 925 Page Mill Road, Palo Alto, CA 94304-1013 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: PURIFIED MANNANASE FROM BACILLUS AMYLOLIQUEFACIENS AND METHOD OF PREPARATION</p> <p>(57) Abstract</p> <p>According to the present invention, a purified mannanase enzyme obtainable from <i>Bacillus amyloliquefaciens</i> is provided. Preferably, the mannanase of the invention has a molecular weight of about 35,000 when analyzed with the SDS-PAGE method and 20,000 when measured by gel filtration, an isoelectric point (pI) of about 5.2-5.6, a pH optimum of about 4.8-5.2 and a half-life at 80 °C of about 45 seconds. In another embodiment of the invention, a method is provided for the preparation of a purified mannanase derived from <i>Bacillus amyloliquefaciens</i>.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CJ	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

PURIFIED MANNANASE FROM *BACILLUS AMYLOLIQUEFACIENS* AND METHOD OF PREPARATION

BACKGROUND OF THE INVENTION

The present invention relates to a novel mannanase enzyme. More specifically, the present invention is related to a novel mannanase obtained from *Bacillus amyloliquefaciens*, and production, purification and use of that mannanase, particularly in the bleaching of pulp.

Industrial uses of hemicellulases have become commercially important in recent years. In addition to animal feed and textiles, pulp and paper applications using hemicellulases continue to expand. While the paper industry has found use for enzymes in pitch control, de-watering and de-inking, these uses are generally still experimental. However, pulp pre-bleaching using xylanase is an established technology. It is believed that xylanase hydrolyzes reprecipitated xylans which are associated with chromophoric lignin on the surfaces of kraft pulp, resulting in improved lignin extractability and bleaching efficiency. With the use of xylanase, consumption of chlorine chemicals is reduced, which significantly reduces levels of hazardous waste released into mill waste streams, halogenated organic compounds being a major by-product of bleaching processes utilizing chlorine and chlorine-containing compounds. As environmental regulations concerning pulp mill effluents increase, chlorine-free alternatives for pulp bleaching will become critical for the paper manufacturing industry. In chlorine-free bleaching sequences, enzymes have been successfully used for increasing brightness or improving pulp quality, in part due to the decreased need for hydrogen peroxide.

Mannan and glucomannans are hemicelluloses associated with xylan and lignin on pulp surfaces and inner layers. As a result, pretreatment of pulp with a combination of mannanase and xylanase has been shown to result in higher brightness than that which can be obtained with xylanase alone (see e.g., Ross et al., *Enzyme-Microb. Technol.*, vol. 14, no. 2, pp. 90-95 (1992)).

Mannanases have been identified in several *Bacillus* organisms. For example, Talbot et al., *Appl. Environ. Microbiol.*, vol. 56, no. 11, pp. 3505-3510 (1990) describes a β -mannanase derived from *Bacillus stearothermophilus* in dimer form having a molecular weight of 162,000 daltons and an optimum pH of 5.5-7.5. Mendoza et. al., *World J. Microbio. Biotech.*, vol. 10, no. 5, pp. 551-555 (1994) describes a β -mannanase derived from *Bacillus subtilis* having a molecular weight of 38,000 daltons, an optimum activity at pH 5.0 and 55 °C and a pI of 4.8. JP 0304706, Derwent Accession No. 91-07734, describes a β -mannanase derived from *Bacillus sp.* having a molecular weight of 37,000 +/- 3,000 daltons when measured by gel filtration, an optimum pH of 8-10 and a pI of 5.3-5.4.

-- 2 --

The prior art, however, fails to identify a mannanase derived from *Bacillus amyloliquefaciens*, as described below.

SUMMARY OF THE INVENTION

According to the present invention, a purified mannanase enzyme obtainable from *Bacillus amyloliquefaciens* is provided. Preferably, the mannanase of the invention has a molecular weight of about 35,000 when analyzed with the SDS-PAGE method and 20,000 when measured by gel filtration, an isoelectric point (pI) of about 5.2-5.6, a pH optimum of about 4.8-5.2 and a half-life at 80 °C of about 45 seconds. In another embodiment of the invention, a method is provided for the preparation of a purified mannanase derived from *Bacillus amyloliquefaciens*.

In another embodiment of the invention, the mannanase of the invention is used in the bleaching of pulp and paper. Preferably, the mannanase is used in combination with a xylanase.

DETAILED DESCRIPTION OF THE INVENTION

According to a preferred embodiment of the invention, a fermentation broth of *Bacillus amyloliquefaciens* is prepared. After separation of whole cells, cell fragments and particulate matter, the supernatant is treated so as to concentrate and isolate the mannanase in the supernatant. In a preferred embodiment of the invention, the purified mannanase comprises a molecular weight of about 33-37 kD as determined with the SDS-PAGE method and about 18-22 kD as determined with the gel filtration method, a pI of about 5.2-5.6, a pH optimum of about 4.8-5.2 and a half life of about 45 seconds at 80 °C. Most preferably, the purified mannanase comprises a molecular weight of about 35 kD as determined with the SDS-PAGE method and about 18-22 kD as determined with the gel filtration method, a pI of about 5.4 and a pH optimum of about 5.0.

Fermentation of *Bacillus amyloliquefaciens* to produce mannanase according to the invention can be accomplished according to any art-recognized method of culturing this microorganism. Preferably, such conditions are manipulated to maximize mannanase production, which conditions are well known in the art.

Purification of the mannanase from the fermentation broth can be by any art-recognized means for purifying such composition. For example, filtration, centrifugation, chromatography, gel filtration or ultrafiltration are all suitable means of purifying the mannanase.

In another embodiment of the invention, the mannanase of the invention is used in the bleaching of pulp for the production of paper. While the use of mannanase alone in the

- 3 -

bleaching of pulp provides an incremental benefit, it is particularly preferred to use mannanase of the invention in combination with xylanase to provide excellent bleaching results. The enzyme treatment (mannanase or mannanase/xylanase) can be utilized at any phase of the bleaching process, however, it is particularly preferred that the mannanase be used prior to bleaching with chlorine-containing chemicals. It is further preferred that the enzyme bleaching be used in combination with oxygen bleaching to limit or even eliminate the use of chlorine-containing chemicals from the bleaching process.

A suitable dosage for mannanase during pulp bleaching is about 10-500 nkat/g pulp when activity is measured by the DNS assay. Conditions for pulp treatment with mannanase are readily ascertained by those of skill in the art, however, a suitable temperature is about 40°C to about 60°C and a suitable pH is from about 4.0 to about 7.0. A suitable dosage for xylanase when used in combination with mannanase is about 0.10 to 200 units/g of dry pulp, and more preferably 0.50 to 50 units/g. The xylanase activity of the enzyme preparations is determined as follows: To 1.8 ml of xylan solution (0.6% Sigma No. X-0627, prepared in 0.05 M sodium acetate buffer and adjusted to pH 5.3 with acetic acid), 0.200 ml of suitably diluted enzyme in the same buffer is added. The solution is incubated at 40°C for exactly 30 minutes. The reaction is then stopped by adding 3 ml DNS reagent (3,5-dinitrosalicylate 10 g/l; Na,K tartrate 300 g/l), and the color is developed by boiling the sample for 5 minutes. The absorbency is then measured at a wave length of 540 nm. One enzyme unit liberates one micromole of reducing sugars calculated at xylose per minute under assay conditions. The activity is calculated from an enzyme dilution liberating 4 micromoles of reducing sugar under assay conditions.

The present invention may be applied to improve any of a wide variety of raw or processed pulps. Processed pulps, i.e., pulps which have been already previously treated to reduce their lignin content, are preferably treated in the process according to the invention to further enhance lignin removal. The present invention is preferably applied to treat softwood pulps to enhance lignin removal and brightening of the pulps. The invention is particularly applicable to chemical pulps, i.e., those in which the lignin component has been chemically modified by various chemical treatments such as in the sulfate (kraft) or sulfite processes and/or oxygen delignification, and is preferably applied to kraft pulps. In a preferred method, the enzymes of the present invention are applied to the pulp after kraft digestion or oxygen delignification but prior to chemical bleaching. In the case where both kraft digestion and oxygen delignification are performed on the same pulp, the enzyme is preferably applied after oxygen delignification. The present invention is also applicable to ozone bleached pulps or pulps which are bleached in ozone containing sequences.

- 4 -

Extractants are often used in pulp bleaching to remove the modified lignin component subsequent to bleaching. In a preferred embodiment of the invention, pulp treated with the enzymes of the present invention is subsequently treated with lignin-degrading chemicals such as chlorine, chlorine dioxide and peroxide, and then with an appropriate extractant. In yet another embodiment, the enzyme treated pulp may be treated with an appropriate extractant, followed by lignin degradation with chemical bleaching or enzymes and a subsequent final treatment with an appropriate extractant. Extractants which solubilize the affected lignin component include bases such as alkali metal hydroxides (E), DMF, dioxane, acetone, and alcohol. Hydroxide extractions may be combined with hydrogen peroxide (E_p) or oxygen (E_o). The resulting pulp may then be further bleached by a chemical bleaching sequence such as chlorine dioxide (DED) or peroxide (P-P) to the desired brightness. Substantial savings of chemicals are observed when the method of the present invention is practiced in comparison with preparing pulp bleached to the same brightness by the same sequence except without using the enzyme treatment by reducing the amount of chlorine-containing chemicals or peroxide used. Similarly, by performing the present invention with the above presented enzymes, one may apply the same amount of bleaching chemicals to the pulp and yet achieve a greater brightness in the treated pulp.

The invention will be further described by reference to the following examples, which examples are illustrative in purpose and are not intended to be limiting.

EXAMPLES

Example 1

Purification of Mannanase from Fermentation Broth of *Bacillus amyloliquefaciens*

Bacillus amyloliquefaciens, ATCC #23842, was grown under conditions including 1% Proflo, 1% locus bean gum and 67.5 ml/l of a medium comprising 8.2 g/l KH_2PO_4 , 91.2 g/l $Na_2HPO_4 \cdot 7H_2O$, 5 g/l $MgSO_4 \cdot 7H_2O$, 10.4 g/l KCl, 11.8 g/l sodium citrate $\cdot 2H_2O$ and 20 g/l yeast extract. The fermentation was carried out for 4-6 days at 37 °C under constant agitation at about 250 rpm in a baffled shake flask. Using a combination of ultrafiltration, gel filtration, and ion exchange chromatography as described below, purified mannanase was obtained. Culture supernatant, 375 ml, enriched for mannanase was concentrated using an Amicon stir-cell (350 ml capacity, PM-10 membrane) to a final volume of 75 ml. 25 ml of this concentrate was applied to a gel filtration column (XK 26/100 packed with Sephacryl S-100 HR, Pharmacia) which had been equilibrated with 10 mM tris-HCl, pH 9.0. The flow rate used was 0.5 ml/min, 15 ml fractions were collected, and UV absorbance was monitored at 280 nm. Mannanase activity in resulting fractions was detected using an RBB-glucomanan substrate assay (see below). Activity eluted in four fractions towards

- 5 -

beginning of main absorbance peak. This procedure was repeated two more times; all mannanase active fractions were pooled. 100 ml of pooled material was then applied to an anion exchange chromatography column (FPLC 10/10 column packed with Q-Sepharose, Pharmacia) equilibrated with 10 mM tris-HCl, pH 9.0. Flow rate was 2 ml/min., UV absorbance was monitored at 280 nm as was conductivity, 1.5 ml fractions were collected. After washing the column with 10 ml (2 void volumes), elution was carried out with a 100 ml linear increasing NaCl gradient, from 10 mM tris-HCl, pH 9.0 to 100 mM NaCl in 10 mM tris-HCl, pH 9.0. Mannanase activity in fractions was determined as above, the majority of which was found to elute approximately mid-way through the salt gradient, in four fractions. Degree of purity was determined using silver stained isoelectric focusing (IEF) gels on a PhastSystem (Pharmacia). IEF gels revealed homogeneous mannanase in the first three of the aforementioned four fractions.

Example 2

Characterization and Properties of Mannanase from *Bacillus amyloliquefaciens*

TECHNIQUES. Relative mannanase activity was determined using a remazol brilliant blue dyed birchwood glucomannan (RBB-mannan) substrate (Megazyme, Sydney, Australia). Samples, 200 μ l, were mixed with 250 μ l of substrate solution (2% (w/v) RBB-mannan in 300 mM sodium acetate pH 4.5) and incubated at 40°C for 10 minutes. Undigested mannan was precipitated by the addition of 1 ml 95% ethanol and removed by centrifugation. Released dye remaining in solution was quantified by spectrophotometry (OD₅₉₀) and was proportional to mannanase activity.

Mannanase activity was quantified using a DNS method for the quantification of resulting reducing sugars. Sample, 200 μ l, was mixed with 1.8 ml galactoglucomannan substrate (0.5% (w/v) locust bean gum in 50 mM sodium citrate, pH 5.3) and incubated for 10 min. at 50 °C. DNS solution (1% (w/v) dinitrosalicylic acid containing 30% (w/v) sodium-potassium tartrate and 1.6% (w/v) NaOH), 3 ml, was added and solution boiled for 5 minutes. OD at 540 nm was measured and was a function of sugar release/mannanase activity as compared to a standard curve. Units are reported in nkat/ml.

Isoelectric focusing (IEF) and sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS PAGE) were carried out using a PhastSystem (Pharmacia) as per manufacturer's instructions. Markers used for pI determination were a broad pI kit pH 3.5-9.3 (Pharmacia Biotech). Molecular weight markers used were from Sigma Chemical Co. (St. Louis, MO). Visualization of proteins was by PhastSystem development silver staining, as per instructions.

Protein concentrations were determined using a BCA method (Pierce Co.).

- 6 -

Molecular weight determinations were accomplished by SDS-PAGE and by gel filtration as follows: Using a Pharmacia FPLC system, 1 ml purified mannanase was applied to two gel filtration columns linked in tandem (Pharmacia Superdex G-200 10/30 followed by Pharmacia Superdex G-75 10/30) which had been equilibrated with 100 mM NaCl-50 mM citrate/phosphate buffer, pH 6.0. Flow rate was 0.5 ml/min., UV absorption was monitored at 280 nm, 1 ml fractions were collected. Fractions were assayed for mannanase activity using an RBB-mannan substrate assay. Mannanase activity was found to elute after 52.5 minutes using this system. Pharmacia low molecular weight gel filtration standards (1.25 mg/ml) were applied to the system using the above conditions and elution results were used to create a molecular weight standard curve. Elution of *Bacillus* mannanase corresponded to a molecular weight of 18-22 kilodaltons when compared to the standard curve.

Thermostability and alkaline stability was determined by adjusting described temperature and pH and assaying as above using the RBB-mannan assay.

A gel overlay method for detecting the presence of multiple mannanases and to determine their isoelectric point (pI) was also developed using RBB-mannan substrate. IEF gels, pH 3-9, were overlaid with a melted agarose-substrate suspension (4% (w/v) agarose, 7 mg/ml RBB-mannan, 0.5% (v/v) glycerol in 50 mM sodium acetate, pH 4.5) and incubated at 37°C. After 1 hour xylanase activity was evident as clearing zones. Gels were allowed to dry completely and stored. Mannanase pI was determined by comparison with identically run IEF gels containing silver stained pI standards.

Table 1
Purified Mannanase Characteristics

	<i>Bacillus amyloliquefaciens</i>
MW (kD, SDS-PAGE)	33-37
MW (kD, Gel Filtration)	18-22
Specific Activity (nkat/mg)	approx. 10500
pI	5.2-5.6
pH Optimum	4.8-5.2
Temp. half life, 80 °C	approx. 45 sec.

Example 3

Pulp Bleaching with Mannanase Based Composition

The bleaching sequence contained the following stages: enzyme treatment, chelation and two alkaline peroxide stages. The enzyme treatment stage was run under the same conditions as described above, except that 10% consistency and culture supernatants enriched for mannanase (purified to be free of cellulase and most xylanase)

- 7 -

were used. The enzyme stage filtrate was acid hydrolyzed and total sugars were detected by HPLC. In the chelation stage, 0.2% EDTA was used at pH 5.5 and 85°C for 30 min at 3% consistency to remove the metals. The peroxide stages were carried out at 10% consistency at 85°C for 4 hours. In the first peroxide stage the H₂O₂ concentration was 3.5% and NaOH concentration 2.2%, in the second the concentrations were 1.5 and 0.85% respectively. After the bleaching stages the pulp was acidified, and handshets were prepared for brightness measurement. The brightness was measured according to the ISO 2469 method. Kappa number, which represents pulp lignin concentration was measured by the SCAN-C1:77 method. Table 2 illustrates the results of pulp bleaching using the XQPP Sequence. "xyl + mann 200" indicates Irgazyme 40 treatment (0.3 l/t) plus mannanase dosed at 200 nkat/g pulp. Monomeric sugars have been detected by HPLC after acid hydrolysis of the enzyme filtrate ("X Stage").

Table 2

Enzyme	X-stage total carbohyd. (g/l)	Brightness ISO %	Kappa
Reference (no enzymes)	0.048	80.4	5.0
Irgazyme Only 0.3 l / t	0.954	81.1	4.4
<i>Bacillus</i> xyl + mann 200	1.548	82.5	4.2
<i>Bacillus</i> : mann 200 only	0.394	80.7	4.9

Of course, it should be understood that a wide range of changes and modifications can be made to the preferred embodiments described above. It is, therefore, intended to be understood that it is the following claims, including all equivalents, which define the scope of the invention.

WE CLAIM

1. A purified mannanase obtained from *Bacillus amyloliquefaciens*.
2. The purified mannanase according to claim 1, wherein said mannanase has a molecular weight of about 33-37 kD as determined by SDS-PAGE.
3. The purified mannanase according to claim 1, wherein said mannanase has a pI of about 5.2-5.6.
4. The purified mannanase according to claim 1, wherein said mannanase has a pH optimum of about 4.8-5.2.
5. The purified mannanase according to claim 1, wherein said mannanase has a half-life of about 45 seconds at a temperature of about 80°C.
6. A method for producing a purified mannanase derived from *Bacillus amyloliquefaciens* comprising:
 - (a) preparing a fermentation broth of *Bacillus amyloliquefaciens*;
 - (b) separating the cells, cell fragments and particulate matter from said fermentation broth to purify said mannanase;
 - (c) optionally concentrating said mannanase to produce a concentrated mannanase solution.
7. The process according to claim 6, wherein said mannanase has a molecular weight of about 33-37 kD as determined by SDS-PAGE.
8. The process according to claim 6, wherein said mannanase has a pI of about 5.2-5.6.
9. The process according to claim 6, wherein said mannanase has a pH optimum of about 4.8-5.2.
10. The process according to claim 6, wherein said mannanase has a half-life of about 45 seconds at a temperature of about 80°C.

- 9 -

11. A process for bleaching pulp comprising contacting said pulp with a composition comprising mannanase derived from *Bacillus amyloliquefaciens*.

12. The process according to claim 11, wherein said composition further comprises xylanase.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 96/14692

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N9/42 D21C9/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N D21C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE WPI Section Ch, Week 9115 Derwent Publications Ltd., London, GB; Class D16, AN 91-105669 XP002022216 "New beta-mannanase hydrolyses beta 1,4-D-mannopyranoside bonds of mannan, glucomannan and can be manufactured at low cost." & JP,A,03 047 076 (AKINO T) , 28 February 1991 cited in the application see abstract	1-3,6-8
Y	---	11,12
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

7 January 1997

Date of mailing of the international search report

15.01.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Mandl, B

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Initial Application No
PCT/US 96/14692

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Section Ch, Week 8816 Derwent Publications Ltd., London, GB; Class D16, AN 88-108330 XP002022217</p> <p>"Production of beta-mannanase which hydrolyses beta-1,4-d-mannopyranoside bonds of e.g. glucomannan etc. and produces manno:oligo:saccharide(s)." & JP,A,63 056 289 (SHINGIJUTSU KAIHATSU, AKINO T) , 10 March 1988 see abstract</p> <p>---</p>	1-3,6-8
X	<p>WO,A,91 18974 (CHEMGEN CORP) 12 December 1991 see page 4, line 8 - line 14; claim 6</p> <p>---</p>	1,6
X	<p>APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 56, no. 11, November 1990, pages 3505-3510, XP000615628 TALBOT G. ET AL.: "Purification and characterization of thermostable beta-mannanase and alpha-galactosidase from Bacillus stearothermophilus." cited in the application see the whole document</p> <p>---</p>	1,6
A		11,12
Y	<p>WO,A,93 24622 (VALTION TEKNILLINEN ;BUCHERT JOHANNA (FI); SIIKA AHO MATTI (FI); V) 9 December 1993 see abstract; example 9; table 5</p> <p>---</p>	11,12
A	<p>US,A,5 296 223 (NAKANISHI KOICHIRO ET AL) 22 March 1994 see the whole document</p> <p>---</p>	1-12
A	<p>ENZYME MICROB. TECHNOL., vol. 14, 2 February 1992, pages 90-95, XP000615629 ROSS N.W. ET AL.: "Enzymic hydrolysis of water-soluble lignin-carbohydrate complexes formed from Populus deltoides: Effects of combinations of beta-mannanases, xylanase and acetyl xylan esterase." see the whole document</p> <p>-----</p>	11,12

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

page 2 of 2

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/US 96/14692

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9118974	12-12-91	AU-A- 8060091	31-12-91
		US-A- 5551515	03-09-96
		US-A- 5476775	19-12-95

WO-A-9324622	09-12-93	AU-A- 4071793	30-12-93
		CA-A- 2135923	09-12-93
		FI-A- 945266	09-11-94
		SE-A- 9404020	23-01-95

US-A-5296223	22-03-94	JP-A- 3167134	19-07-91
		CA-A- 2030770	25-05-91
		DE-D- 69009706	14-07-94
		DE-T- 69009706	29-09-94
		EP-A- 0431811	12-06-91

Form PCT/ISA/210 (patent family annex) (July 1992)